arXiv:1612.09016v1 [math.AG] 29 Dec 2016

A CRITERION FOR THE PRIMITIVITY OF A BIRATIONAL AUTOMORPHISM OF A CALABI-YAU MANIFOLD AND AN APPLICATION

KEIJI OGUISO

Abstract. We shall give a sufficient condition on the primitivity of a birational automrphism of a Calabi-Yau manifold in purely algebro geometric terms. As an application, we shall give an explicit construction of Calabi-Yau manifolds of Picard number 2 of any dimension ≥ 3, with primitive birational automrphisms of first dynamical degree > 1.

1. Introduction

Throughout this note, we work in the category of projective varieties defined over $\mathbb{C}$. This note is a continuation of [Og14], [Og16-2] and is much inspired by recent works of Bianco [Bi16] in technique and also works of Amerik-Verbitsky [AV16] and Ouchi [Ou16] in spirit for dynamical studies of automorphisms of hyperkähler manifolds.

Our main interest is the existence of primitive birational automorphisms on Calabi-Yau manifolds in any dimension ≥ 3, especially of the smallest possible Picard number 2 as done by [AV16], [Ou16] for hyperkähler manifolds.

Definition 1.1. Let V be a projective variety and $f \in \mathrm{Bir}(V)$. We call a rational dominant map $\pi : V \dashrightarrow B$ to a projective variety B with connected fibers a *rational fibration*. f is said to be *non-trivial* if $0 < \dim B < \dim V$. A rational fibration $\pi : V \dashrightarrow B$ is said to be *f-equivariant* if there is $f_B \in \mathrm{Bir}(B)$ such that $\pi \circ f = f_B \circ \pi$. We say that f is *primitive* if there is no non-trivial f-equivariant rational fibration $\pi : V \dashrightarrow B$.

The notion of primitivity is introduced by De-Qi Zhang [Zh09]. Primitivity of a birational automorphism in the category of projective varieties is an analogue of irreducibility of a linear selfmap in the category of finite dimensional linear spaces.

Our aim is to prove the following two theorems (Theorems 1.2, 1.3; see also Remark 1.4 for notions in the theorems):

Theorem 1.2. *Let X be a minimal Calabi-Yau variety of dimension $n \geq 3$ of Picard number $\rho(X) \geq 2$ and $f \in \mathrm{Bir}(X)$ such that*

1. *For any movable divisor D, there are a minimal Calabi-Yau variety X' and a birational map $g : X' \dashrightarrow X$, both allowing to depend on D, such that $D' = g^*D$ is semi-ample on X'; and*
2. *The action $f^*|N^1(X)_{\mathbb{Q}}$ is irreducible over $\mathbb{Q}$.*

Then f is primitive. If in addition that X is smooth, then $d_1(f) > 1$ as well.

The author is supported by JSPS Grant-in-Aid (S) No 25220701, JSPS Grant-in-Aid (S) 15H05738, JSPS Grant-in-Aid (A) 16H02141, JSPS Grant-in-Aid (B) 15H03611, and by KIAS Scholar Program.

Theorem 1.3. *For each $n \geq 3$, there is an n-dimensional Calabi-Yau manifold M of Picard number $\rho(M) = 2$ with a primitive birational automorphism $f \in \mathrm{Bir}(M)$ of the first dynamical degree $d_1(f) > 1$.*

Our manifold M and $f \in \mathrm{Bir}(M)$ in Theorem 1.3 are explicit (See Section 3). If f is primitive, then $\mathrm{ord}(f) = \infty$ ([Og16-1, Lemma 3.2]). In particular, $\rho(M) \geq 2$ if M is a Calabi-Yau manifold with primitive $f \in \mathrm{Bir}(M)$ (see eg. [Og14]). So, Theorem 1.3 also shows that the estimate $\rho(M) \geq 2$ is optimal for Calabi-Yau manifolds with primitive birational automorphisms in each dimension ≥ 3.

Theorem 1.2 may have its own interest, suggests some relation of primitivity of birational automorphisms and primitivity (= irreducibility) of its geometrically meaningful linear representation. The condition (1) in Theorem 1.2 is automatical if the log minimal model program works in dimension n (then one makes D' nef) and log abundance theorem also holds in dimension n (then D' is semi-ample). Theorem 1.2 is also crucial to prove Theorem 1.3. We prove Theorem 1.2 in Section 2.

We refer [Og15], [Og16-2] and references therein for background and known results relevant to our main theorems. For convenience to the readers, we recall some definitions and facts, which we need:

Remark 1.4. Let X be a normal projective variety of dimension $\ell \geq 2$ and $f \in \mathrm{Bir}(X)$.

(1) We call X a *Calabi-Yau manifold*, if X is smooth, simply-connected, $H^0(\Omega_X^j) = 0$ for $0 < j < \ell$ and $H^0(\Omega_X^\ell) = \mathbb{C}\omega_X$ for a nowhere vanishing regular ℓ-form ω_X. We call X a *minimal Calabi-Yau variety* if X has at most $\mathbb{Q}$-factorial terminal singularities, $h^1(\mathcal{O}_X) = 0$ and $\mathcal{O}_X(K_X) \simeq \mathcal{O}_X$.

(2) Following [Ka88], we say that a $\mathbb{Q}$-Cartier Weil divisor D is *movable* (resp. *semi-ample*) if there is a positive integer m such that the complete linear system $|mD|$ has no fixed component (resp. is free).

(3) For a rational map $g : X \dashrightarrow Y$ from a smooth projective variety X to a smooth projective variety Y, we define $g^* : N^p(Y) \to N^p(X)$ by

$$g^* = (p_1)_* \circ p_2^* .$$

Here $N^p(X)$ is a free $\mathbb{Z}$-module generated by the numerical equivalence classes of p-cocycles, $p_1 : Z \to X$ is a Hironaka resolution ([Hi64]) of the indeterminacy of g, $p_2 : Z \to Y$ is the induced morphism, p_2^* is the pullback as a cocycle and $(p_1)_*$ is the pushforward as a cycle ([Fu84, Chap19, Example 19.1.6]). In this definition, smoothness of Y is not needed but smoothness of Z and X are needed in order to identify cycles and cocycles over $\mathbb{Z}$.

(4) Assume that X is smooth. The p-th *dynamical degree* of $f \in \mathrm{Bir}(X)$ is defined by

$$d_p(f) := \lim_{n\to\infty} (((f^n)^* H^p).H^{\ell-p})_X^{\frac{1}{n}} .$$

Here H is any ample divisor on X. By Dinh-Sibony [DS05] (see also [Tr15]), $d_p(f)$ is well-defined, independent of the chioce H, and birational invariant in the sense that $d_p(\varphi^{-1} \circ f \circ \varphi) = d_p(f)$ for any birational map $\varphi : X' \dashrightarrow X$ from a smooth projective variety X'.

(5) Assume that X is not necessarily smooth but $\mathbb{Q}$-factorial. Then any Weil divisors (codimension one cycles) are $\mathbb{Q}$-Cartier divisors (codimension one cocycles) on X. We denote by $N^1(X)_{\mathbb{Q}}$ the finite dimensional $\mathbb{Q}$-linear space spanned by the numerical equivalence classes of Weil divisors on X over $\mathbb{Q}$. So, in this case, one can define the map $g^* : N^1(Y)_{\mathbb{Q}} \to N^1(X)_{\mathbb{Q}}$ for a rational map $g : X \dashrightarrow Y$ in the same way as in (3) (over $\mathbb{Q}$, not necessarily over $\mathbb{Z}$).

(6) Assume that either X is $\mathbb{Q}$-factorial and $f \in \mathrm{Bir}\,(X)$ is isomorphic in codimension one, for instance, this is the case when X is a minimal Calabi-Yau variety (see eg. [Ka08, Page 420]), or $f \in \mathrm{Aut}\,(X)$. Then, $d_1(f)$ is well-defined by the same formula as in (4) using (5). Moreover, the correspondence $f \mapsto f^*$ is functorial in the sense that $(f \circ g)^* = g^* \circ f^*$ for $f, g \in \mathrm{Bir}\,(X)$ in the first case and for any $f, g \in \mathrm{Aut}\,(X)$ in the second case. In particular, $f^* \in \mathrm{GL}\,(N^1(X)_{\mathbb{Q}})$. Moreover, $d_1(f)$ coincides with the spectral radius of $f^*|N^1(X)_{\mathbb{Q}}$ by $(f^n)^* = (f^*)^n$ for any $n \in \mathbb{Z}$. One observes this by writing the Jordan canonical form of $f^*|N^1(X)_{\mathbb{Q}}$ over $\mathbb{C}$.

Acknowledgement. I would like to express my thanks to Professors Ekaterina Amerik, Ljudmila Kamenova and De-Qi Zhang for their interest in this work and useful discussions.

2. Proof of Theorem 1.2.

In this section, we prove Theorem 1.2.

Lemma 2.1. *X has no f-equivariant rational fibration $\pi : X \dashrightarrow B$ such that $0 < \dim B < \dim X$ and $\kappa(\tilde{X}_b) = 0$ for general $b \in B$. Here $\nu : \tilde{X} \to X$ is a Hironaka resolution of the indeterminacy of π and the singularities of X ([Hi64]), $\tilde{X}_b$ is the fiber over $b \in B$ of the morphism $\tilde{\pi} := \pi \circ \nu : \tilde{X} \to B$ and $\kappa(\tilde{X}_b)$ is the Kodaira dimension of $\tilde{X}_b$.*

Proof. Assuming to the contray that X has an f-equivariant rational fibration $\pi : X \dashrightarrow B$ such that $0 < \dim B < \dim X$ and $\kappa(\tilde{X}_b) = 0$ for general $b \in B$, we shall derive a contradiction.

By taking a Hironaka resolution of singularities, we may and will assume that B is smooth. Let H be a very ample divisor on B. Set $L := \nu_*(\tilde{\pi}^* H)$. Then $|L|$ is movable and $\pi = \Phi_\Lambda$. Here $\Lambda \subset |L|$ is a sublinear system of $|L|$ and Φ is the rational map associated to Λ. Then, by the assumption (1) in Theorem 1.2, there are a minimal Calabi-Yau variety X' and a birational map $g : X' \dashrightarrow X$ such that g^*L is semi-ample. So, by replacing (X, f) by

$$(X', f' := g^{-1} \circ f \circ g) \ ,$$

we may and will assume that $|L|$ is semi-ample. Note that $(f')^*|N^1(X')_{\mathbb{Q}}$ is irreducible over $\mathbb{Q}$, as $g^{\pm 1}$ and f are isomorphic in codimension one so that

$$(f')^* = (g^{-1} \circ f \circ g)^* = g^* \circ f^* \circ (g^*)^{-1} \ .$$

Let us take a sufficiently large positive integer m such that $|mL|$ is free and the morphism $\varphi_m = \Phi_{|mL|}$ is the Iitaka-Kodaira fibration associated to L. Set $B' = \varphi(X)$. Then $\dim B' = \kappa(L, X)$. Here $\kappa(L, X)$ is the Iitaka-Kodaira dimension of L. As $\Lambda \subset |L| \subset |mL|$, the surjective morphism $\varphi_m : X \to B'$ factors through $\pi : X \dashrightarrow B$, i.e., there is a (necessarily dominant) rational map $\rho : B' \dashrightarrow B$ such that $\pi = \rho \circ \varphi_m$.

Claim 2.2. $\rho : B' \dashrightarrow B$ is a birational map.

Proof. As π is of connected fibers and ρ is dominant, it suffices to show that $\dim B = \dim B'$, i.e., that $\dim B = \kappa(L, X)$. Set $\tilde{L} := \nu^* L$. Then $\kappa(L, X) = \kappa(\tilde{L}, \tilde{X})$. In what follows, we shall prove $\dim B = \kappa(\tilde{L}, \tilde{X})$.

Let $\{E_i\}_{i=1}^m$ be the set of exceptional prime divisors of ν. Then we can write

$$\nu^* L = \tilde{\pi}^* H + \sum_{i=1}^m a_i E_i \ , \ K_{\tilde{X}} = \sum_{i=1}^m b_i E_i \ .$$

Here $a_i \geq 0$ as ν resolves the indeterminacy of π and $b_i > 0$ as X is a minimal Calabi-Yau variety. Note that $\tilde{L}|_{\tilde{X}_b} = a_i E_i|_{\tilde{X}_b}$ as $H|_{\tilde{X}_b}$ is trivial and $K_{\tilde{X}_b} = b_i E_i|_{\tilde{X}_b}$ by the adjunction formula. Then there is a positive integer m such that $mK_{\tilde{X}_b} - \tilde{L}|_{\tilde{X}_b}$ is linearly equivalent to an effective divisor. Hence

$$0 \leq \kappa(\tilde{L}|_{\tilde{X}_b}, \tilde{X}_b) \leq \kappa(\tilde{X}_b) = 0 \ ,$$

as $b \in B$ is general and $\kappa(\tilde{X}_b) = 0$ by the assumption. Thus $\kappa(\tilde{L}|_{\tilde{X}_b}, \tilde{X}_b) = 0$. Therefore

$$\dim B \leq \kappa(\tilde{L}, \tilde{X}) \leq \dim B + \kappa(\tilde{L}|_{\tilde{X}_b}, \tilde{X}_b) = \dim B \ .$$

Here the second inequality follows from [Ue75, Theorem 5.11]. Hence $\dim B = \kappa(\tilde{L}, \tilde{X})$ as desired. □

By Claim 2.2, we may and will assume that $\pi : X \to B$ is an f-equivariant surjective *morphism* given by the free complete linear system $|L|$, by replacing $\pi : X \dashrightarrow B$ by $\varphi_m : X \to B'$ for sufficiently large divisible m. Then $B = \mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kL))$ and $\pi = \Phi_{|L|}$.

Claim 2.3. There is $f_B \in \mathrm{Aut}\,(B)$ (*not only in* $\mathrm{Bir}\,(B)$) such that $f_B \circ \pi = \pi \circ f$ as a rational map from X to B.

Proof. As f is isomorphic in codimension one, the pullback f^* induces an isomorphism

$$f_* : B = \mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kL)) \simeq \mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kf^*L)) = B \ .$$

Here the last equality is the one under the identification of $\mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kL))$ with $\mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kf^*L))$ defined by $D \mapsto f^*D$ for $D \in |kL|$.

Then the image of $\pi \circ f = \Phi_{|f^*L|}$ is $B = \mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kf^*L))$ under the last identification made above. Thus

$$f_* \in \mathrm{Aut}\,(B)$$

under the first identification $B = \mathrm{Proj} \oplus_{k \geq 0} H^0(X, \mathcal{O}_X(kL))$ and f_* satisfies that $\pi \circ f = f_* \circ \pi$. We may now take $f_B = f_*$. □

As $f_B \in \mathrm{Aut}\,(B)$, the map $f_B^* : N^1(B)_{\mathbb{Q}} \to N^1(B)_{\mathbb{Q}}$ is a well-defined isomorphism (cf. Remark 1.4). Note also that

$$\pi^* \circ f_B^* = f^* \circ \pi^* \ .$$

Indeed, $(f_B \circ \pi)^* = \pi^* \circ f_B^*$, as π and f_B are morphisms. We have also $(\pi \circ f)^* = f^* \circ \pi^*$, as f is isomorphic in codimension one and X is normal and $\mathbb{Q}$-factorial (cf. Remark 1.4). Thus $\pi^* \circ f_B^* = f^* \circ \pi^*$ from $f_B \circ \pi = \pi \circ f$. Hence the subspace $\pi^* N^1(B)_{\mathbb{Q}} \subset N^1(X)_{\mathbb{Q}}$ is

f-stable by $f_B^*(N^1(B)_{\mathbb{Q}}) = N^1(B)_{\mathbb{Q}}$. On the other hand, as $0 < \dim B < \dim X$, X and B are projective and π is a *morphism*, it follows that

$$0 \neq \pi^* N^1(B)_{\mathbb{Q}} \neq N^1(X)_{\mathbb{Q}} \ ,$$

a contradiction to the irreducibility of the action f^* on $N^1(X)_{\mathbb{Q}}$, the assumption (2) in Theorem 1.2. This completes the proof of Lemma 2.1. □

Lemma 2.4. *Let P be a very general closed point of X. Then f^n is defined at P for all $n \in \mathbb{Z}$ and the set $\{f^n(P) \,|\, n \in \mathbb{Z}\}$ is Zariski dense in X.*

Proof. The first assertion is clear. We show the second assertion. By [AC13, Théorème 4.1], there is a smooth projective variety C and a dominant rational map $\rho : X \dashrightarrow C$ such that $\rho \circ f = \rho$ as a rational map and $\rho^{-1}(\rho(P))$ is the Zariski closure of $\{f^n(P) \,|\, n \in \mathbb{Z}\}$ for very general $P \in X$.

Recall that the map $\rho^* : N^1(C)_{\mathbb{Q}} \to N^1(X)_{\mathbb{Q}}$ is well-defined, as X is normal and $\mathbb{Q}$-factorial (Remark 1.4). As f is isomorphic in codimension one and $\rho = \rho \circ f$, we have

$$\rho^* = f^* \circ \rho^* : N^1(C)_{\mathbb{Q}} \to N^1(X)_{\mathbb{Q}}$$

as in the proof of Lemma 2.1 (See also [Bi16, Lemma 4.5]). Thus

$$f^* | \rho^* N^1(C)_{\mathbb{Q}} = id_{\rho^* N^1(C)_{\mathbb{Q}}} \ .$$

As $\rho(X) \geq 2$ and f is irreducible on $N^1(X)_{\mathbb{Q}}$ by our assumptions in Theorem 1.2, it follows that

$$\rho^* N^1(C)_{\mathbb{Q}} = \{0\} \ .$$

As C is projective, this is possible only when $\dim C = 0$, i.e., C is a point. This implies the second assertion. □

The following important proposition is proved by Bianco [Bi16, Proposition (3) and Section 4]:

Proposition 2.5. *Let X be a projective variety (not necessarily a minimal Calabi-Yau variety) and $f \in \mathrm{Bir}(X)$. Assume that $\pi : X \dashrightarrow B$ is a nontrivial f-equivariant rational fibration such that $\tilde{X}_b$ is of general type for general $b \in B$. Here $\nu : \tilde{X} \to X$ is a Hironaka resolution of the singularities of X and the indeterminacy of π and $\tilde{X}_b$ is the fiber over $b \in B$ of the morphism $\tilde{\pi} := \pi \circ \nu : \tilde{X} \to B$. Then for any very general point $P \in X$ at which f^n ($n \in \mathbb{Z}$) are defined, the well-defined $\langle f \rangle$-orbit $\{f^n(P) | n \in \mathbb{Z}\}$ is never Zariski dense in X.*

Remark 2.6. The assertion [Bi16, Proposition (2)] seems a bit too optimistic, and the proof of [Bi16, Proposition (3)], which we cited above, uses [Bi16, Proposition (2)]. However, one may avoid the use of [Bi16, Proposition (2)] by the following slight modification of the original proof of [Bi16, Proposition (3)].

By taking a Hironaka resolution of B, X and the indeterminacy $X \dashrightarrow B$, we may assume that $\pi : X \to B$ is a surjective morphism between smooth projective varieties. We may also assume that for any very general point $Q \in B$ at which f_B^n ($n \in \mathbb{Z}$) are defined, the well-defined $\langle f_B \rangle$-orbit $\{f_B^n(Q) | n \in \mathbb{Z}\}$ is Zariski dense in B; as otherwise, the assetion is obvious.

First we take a Zariski dense open subset $U \subset B$ such that $\pi_U := \pi|\pi^{-1}(U) : \pi^{-1}(U) \to U$ is a smooth morphism. Next we take the relative canonical model

$$\pi_U^{\mathrm{can}} : Y := \mathrm{Proj} \oplus_{m\geq 0} (\pi_U)_* \mathcal{O}_X(mK_{\pi^{-1}(U)}) \to U$$

of $\pi_U : \pi^{-1}(U) \to U$ over U (see [BCHM10, Corollary 1.1.2], [Ka09, Theorem 6.6] for the finite generation). Then, as in [Bi16, Proof of Proposition (1)], $\pi_U^{\mathrm{can}} : Y \to U$ is isotrivial, by the Zariski density of $\langle f_B \rangle$-orbit. In particular, possibly after shrinking U a bit, all fibers Y_b ($\in U$) of π_U^{can} are isomorphic, say to F, which is the canonical model of the fibers. Then, by [Ue75, Corollary 14.3], $\mathrm{Bir}\,(F) = \mathrm{Aut}\,(F)$ and this group is a finite group, say of order k, and the induced birational map $f_b^n : Y_b \to Y_{f_B^n(b)}$ is an isomorphism for $b \in U$ at which the rational map $f_B^n|U : U \dashrightarrow U$ is defined. Moreover, by the isotriviality, there is an open covering

$$U = \cup_{i\in I} U_i$$

of U in the classical topology such that $Y \to U$ is isomorphic to $F \times U_i$ over U_i ([Se06, Proposition 2.6.10]). As $\mathrm{Aut}\,(F)$ is finite, local trivial pieces $F \times U_i$ and $F \times U_j$ are glued by a *constant* $g_{ij} \in \mathrm{Aut}\,(F)$ over $U_i \cap U_j$ to be $Y \to U$. As $P \in X$ is a very general point, we may identify $P \in X$ with the unique corresponding point in Y, which we denote by the same letter P. Note then that the isomorphism above $f_b^n : Y_b \to Y_{f_B^n(b)}$ is also represented by $\mathrm{Aut}\,(F)$ if we choose a local chart $F \times U_i$ and $F \times U_j$ such that $b \in U_i$ and $f_B^n(b) \in U_j$. Hence the $\langle f \rangle$-orbit of $P \in Y$ is contained in a multisection $\tilde{U} \ni P$ of $Y \to U$ such that $P \in \tilde{U} \in Y$ and of degree at most k over U. This implies Proposition 2.5.

Now we are ready to complete the proof of Theorem 1.2.

Assume that X admits a non-trivial f-equivariant rational fibration $\pi : X \dashrightarrow B$. Let $\tilde{\pi} : \tilde{X} \to B$ be a Hironaka resolution of indeterminacy of π and $\mathrm{Sing}\,(X)$. Consider the relative Kodaira fibration over B (see [BCHM10, Corollary 1.1.2], [Ka09, Theorem 6.6] for the finite generation):

$$g : X \dashrightarrow K := \mathrm{Proj} \oplus_{m\geq 0} \tilde{\pi}_* \mathcal{O}_{\tilde{X}}(mK_{\tilde{X}}) \ .$$

Then g is f-equivariant and $\kappa(\tilde{X}_k) = 0$ for general $k \in K$. Here $\tilde{X}_k$ is a Hironaka resolution of the fiber over $k \in K$. By Lemma 2.4 and Proposition 2.5 and by $\dim B > 0$, we have $0 < \dim K < \dim X$. However, this contradicts to Lemma 2.1. Thus f is primitive.

We shall show that $d_1(f) > 1$ if X is smooth. As f is isomorphic in codimension one, f^* preserves the movable cone $\overline{\mathrm{Mov}}(X) \subset N^1(X)_{\mathbb{R}}$, which is, by definition, the closed convex hull of the movable divisor classes in $N^1(X)_{\mathbb{R}}$. As $\overline{\mathrm{Mov}}(X)$ is a strictly convex closed cone in $N^1(X)_{\mathbb{R}}$, it follows from the Birkhoff-Perron-Frobenius theorem ([Br67]) that there is $0 \neq v \in \overline{\mathrm{Mov}}(X)$ such that $f^*v = d_1(f)v$. As X is smooth (hence the Weil divisors are Cartier divisors), $f^* : N^1(X) \to N^1(X)$ is a well-defined isomorphism over $\mathbb{Z}$ (not only over $\mathbb{Q}$). Thus the product of the eigenvalues of f^* is of absolute value 1, as $\mathbb{Z}^\times = \{\pm 1\}$. Thus $d_1(f) \geq 1$. If $d_1(f) = 1$, then there would be $0 \neq u \in N^1(X)_{\mathbb{Q}}$ such that $f^*(u) = u$. Hence $d_1(f) > 1$, as $f^*|N^1(X)_{\mathbb{Q}}$ is irreducible over $\mathbb{Q}$ and $\rho(X) \geq 2$ by our assumptions in Theorem 1.2. This completes the proof of Theorem 1.2.

3. Proof of Theorem 1.3.

In this section, we shall prove Theorem 3.1 below, from which Theorem 1.3 follows.

Calabi-Yau manifolds in Theorem 3.1 are higher dimensional generalization of Calabi-Yau threefolds studied in [Og14, Section 6].

Let n be an integer such that $n \geq 3$. Let

$$M = H_1 \cap H_2 \cap \ldots \cap H_{n-1} \cap Q \subset \mathbb{P}^n \times \mathbb{P}^n$$

be a general complete intersection of $n-1$ hypersurfaces H_i $(1 \leq i \leq n-1)$ of bidegree $(1,1)$ and a hypersurface Q of bidegree $(2,2)$ in $\mathbb{P}^n \times \mathbb{P}^n$. Then, by the Lefschetz hyperplane section theorem, M is a smooth Calabi-Yau manifold of dimension n and of Picard number 2. More precisely,

$$\mathrm{Pic}\,(M) \simeq N^1(M) = \mathbb{Z}h_1 \oplus \mathbb{Z}h_2 \ .$$

Here and hereafter, $p_i : \mathbb{P}^n \times \mathbb{P}^n \to \mathbb{P}^n$ is the projection to the i-th factor, L_i is the hyperplane class of the i-th $\mathbb{P}^n$, $H_i := p_i^* L_i$, $h_i = H_i|_M$ $(i = 1, 2)$.

Let

$$V = \cap_{i=1}^{n-1} H_i \subset \mathbb{P}^n \times \mathbb{P}^n \ .$$

Then V is a smooth Fano manifold of dimension $n+1$ with $\mathrm{Pic}\,(V) = \mathbb{Z}H_1|V \oplus \mathbb{Z}H_2|V$ and $M \in |-K_V|$. Note that $H_i|V$ gives the i-th projection $p_i|V : V \to \mathbb{P}^n$ to the i-th factor. In particular, $H_i|V$ is free (hence nef) but not ample. Then, by a result of Kollár [Bo91, Appendix], we have $\overline{\mathrm{Amp}}\,(V) \simeq \overline{\mathrm{Amp}}\,(M)$ under the inclusion map $M \subset V$, and therefore

$$\overline{\mathrm{Amp}}\,(M) = \mathbb{R}_{\geq 0}h_1 + \mathbb{R}_{\geq 0}h_2$$

in $N^1(M)_{\mathbb{R}} := N^1(M) \otimes_{\mathbb{Z}} \mathbb{R}$. In particular, any nef divisor on M is semi-ample.

We also note that $g^*(h_1 + h_2) = h_1 + h_2$ for $g \in \mathrm{Aut}\,(M)$. Hence $|\mathrm{Aut}\,(M)| < \infty$ (see eg. [Og14, Proposition 2.4]). So, if $f \in \mathrm{Bir}\,(M)$ is primitive, then necessarily $f \notin \mathrm{Aut}\,(M)$ in our case, as f has to be of infinite order ([Og16-1, Lemma 3.2]).

Consider the projections

$$\pi_i := p_i|M : M \to \mathbb{P}^n$$

$(i = 1, 2)$. Then π_1 are of degree 2 by the definition of M. Hence we have a birational involutions $\tau_i \in \mathrm{Bir}\,(M)$ corresponding to p_i.

Our main result of this section is the following:

Theorem 3.1. *Under the notation above, $f := \tau_1 \circ \tau_2$ is a primitive birational automorphism of M with*

$$d_1(f) = (2n^2 - 1) + 2n\sqrt{n^2 - 1} > 1 \ .$$

Proof. In the proof, we will frequently use the fact that any birational automorphism of M is isomorphic in codimension one (Remark 1.4).

Lemma 3.2. *With respect to the basis $\langle h_1, h_2 \rangle$, the actions of $\tau_i^*|N^1(M)$ $(i = 1, 2)$ and $f^*|N^1(X)$ are represented by the following matrices M_i, M_2M_1 respectively:*

$$M_1 = \begin{pmatrix} 1 & 2n \\ 0 & -1 \end{pmatrix} \ , \ M_2 = \begin{pmatrix} -1 & 0 \\ 2n & 1 \end{pmatrix} \ , \ M_2M_1 = \begin{pmatrix} -1 & -2n \\ 2n & 4n^2 - 1 \end{pmatrix}.$$

The eigenvalues of f^* are

$$(2n^2 - 1) \pm 2n\sqrt{n^2 - 1} \ ,$$

which are irrational. In particular, $f^*|N^1(X)_{\mathbb{Q}}$ is irreducible over $\mathbb{Q}$.

Proof. We have $\tau_1^* h_1 = h_1$. We can write $(\pi_1)_* h_2 = aL_1$ for some integer a. First, we detemine the value a. As $h_2 = \pi_2^* L_2$, we readily compute that

$$a = (aL_1.L^{n-1})_{\mathbb{P}^n} = ((\pi_1)_* \pi_2^* L_2 . L_1^{n-1})_{\mathbb{P}^n} = (\pi_2^* L_2 . \pi_1^* L_1^{n-1})_M$$
$$= (H_2 . H_1^{n-1} . 2(H_1 + H_2)^n)_{\mathbb{P}^n \times \mathbb{P}^n} = 2n \ .$$

Thus

$$h_2 + \tau_1^* h_2 = p_1^*(p_1)_* h_2 = 2nh_1 \ .$$

Hence $\tau_1^* h_1 = h_1$ and $\tau_2^* h_2 = 2nh_1 - h_2$, and therefore, the matrix representaion M_1 of τ_1^* is as described. In the same way, one obtains the matrix representation M_2 of τ_2^* as described. The matrix representaion of f^* is then $M_2 M_1$. The rest follows from a simple computaion of 2×2 matrices and an elementary fact that $\sqrt{n^2 - 1}$ is irrational for any integer $n \geq 2$. □

Let $\tilde{\pi}_i : M \to M_i$ $(i = 1, 2)$ be the Stein factorization of $\pi_i : M \to \mathbb{P}^n$. As the Stein factorization is unique, the covering involution $\tilde{\tau}_i$ of $M_i \to \mathbb{P}^n$ is in $\mathrm{Aut}\,(M_i)$ (not only in $\mathrm{Bir}\,(M_i)$) and satisfies $\tilde{\pi}_i \circ \tau_i = \tilde{\tau}_i \circ \tilde{\pi}_i$.

Lemma 3.3. *$\tilde{\pi}_i$ $(i = 1, 2)$ are small contractions of M and M admits no other contraction. Here a contarction of M means a non-isomorphic surjective morphism to a normal projective variety of positive dimension with connected fibers.*

Proof. The morphism $\tilde{\pi}_i$ is given by $|mh_i|$ for large m. Recall that h_i are not ample. Thus $\tilde{\pi}_i$ is a contraction. As $\rho(M) = 2$, there is then no contraction other than $\tilde{\pi}_i$ $(i = 1, 2)$.

As $\rho(V) = 2$, the i-th projection $p_i|V : V \to \mathbb{P}^n$ from V $(i = 1, 2)$ contracts no divisor to a subvariety of codimension ≥ 2. Thus, the i-th projection $\pi_i = p_i|M : M \to \mathbb{P}^n$ contracts no divisor, as $M = V \cap Q$ and Q is a general very ample divisor on V. Hence $\tilde{\pi}_i$ is a small contraction. The fact that $\tilde{\pi}_i$ $(i = 1, 2)$ are small contractions of M also follows from the proof of the next Lemma 3.4. □

Lemma 3.4. $\mathrm{Bir}\,(M) = \mathrm{Aut}\,(M) \cdot \langle \tau_1, \tau_2 \rangle$.

Proof. As $\rho(M) = 2$, the relative Picard number $\rho(M/M_1)$ is 1. By Lemma (3.2), we have

$$\tau_1^* h_2 = -h_2 + 2nh_1 \ .$$

Thus $\tau_1^* h_2$ is relatively anti-ample for $\overline{\tau}_1^{-1} \circ \tilde{\pi}_1 : M \to M_1$, while h_2 is relatively ample for $\tilde{\pi}_1 : M \to M_1$. Since $K_M = 0$, the map $\tilde{\tau}_1^{-1} \circ \tilde{\pi}_1 : M \to M_1$ is then the flop of $\tilde{\pi}_1 : X \to M_1$, given by τ_1. For the same reason, the map $\tilde{\tau}_2^{-1} \circ \tilde{\pi}_2 : M \to M_2$ is the flop of $\tilde{\pi}_2 : M \to M_2$, given by τ_2.

Recall that any flopping contraction of a Calabi-Yau manifold is given by a codimension one face of $\overline{\mathrm{Amp}}\,(M)$ up to automorphisms of M ([Ka88, Theorem 5.7]). As there is no codimension one face of $\overline{\mathrm{Amp}}\,(M)$ other than $\mathbf{R}_{\geq 0} h_i$ $(i = 1, 2)$, there is then no flop of M other than τ_i $(i = 1, 2)$ up to $\mathrm{Aut}\,(M)$. Recall a fundamental result of Kawamata ([Ka08, Theorem 1]) that *any birational map between minimal models is decomposed into finitely many flops up to automorphisms of the target variety.* Thus any $\varphi \in \mathrm{Bir}\,(M)$ is decomposed into a finite sequence of flops τ_i and an automorphism of M at the last stage. This proves the result. □

Set $v_+ := -h_1 + (n + \sqrt{n^2-1})h_2$, $v_- := -h_2 + (n + \sqrt{n^2-1})h_1$ and

$$V := \mathbb{R}_{\geq 0}v_+ + \mathbb{R}_{\geq 0}v_- \subset N^1(M)_{\mathbb{R}} \ .$$

Here $v_\pm$ are eigenvectors of f^* corresponding to the eigenvalues $(2n^2-1) \pm 2n\sqrt{n^2-1}$. By writing the Jordan canonical form of the matrix M_2M_1 in Lemma 3.2, one readily observes that

$$\lim_{n\to\infty} \mathbb{R}_{>0}(f^n)^*x = \mathbb{R}v_+ \ , \ \lim_{n\to\infty} \mathbb{R}_{>0}(f^{-n})^*x = \mathbb{R}v_- \ --- (*)$$

for any $x \in \overline{\mathrm{Amp}}\,(M) \setminus \{0\}$.

Lemma 3.5. $\overline{\mathrm{Mov}}\,(M) = V$. *Moreover, the interior* $(\overline{\mathrm{Mov}}\,(M))^\circ$ *of* $\overline{\mathrm{Mov}}\,(M)$ *coincides with the* $\mathrm{Bir}\,(M)$*-orbit of the nef cone* $\overline{\mathrm{Amp}}\,(M)$*:*

$$\overline{\mathrm{Mov}}\,(M)^\circ = \mathrm{Bir}\,(M)^*\overline{\mathrm{Amp}}\,(M) := \cup_{h\in \mathrm{Bir}\,(M)} h^*\overline{\mathrm{Amp}}\,(M) \ .$$

Proof. Recall that $\mathrm{Aut}\,(M)$ preserves $\overline{\mathrm{Amp}}\,(M) = \mathbb{R}_{\geq 0}h_1 + \mathbb{R}_{\geq 0}h_2$, in particular,

$$g^*\{h_1, h_2\} = \{h_1, h_2\}$$

if $g \in \mathrm{Aut}\,(M)$ and $\mathrm{Bir}\,(M)$ preserves $\overline{\mathrm{Mov}}\,(M)$. As every nef divisor on M is semi-ample in our situation, we have

$$\mathrm{Bir}\,(M)^*\overline{\mathrm{Amp}}\,(M) \subset \overline{\mathrm{Mov}}\,(M) \ .$$

By using (*), we also find that

$$V^\circ = \mathrm{Bir}\,(M)^*\overline{\mathrm{Amp}}\,(M)$$

and therefore

$$V \subset \overline{\mathrm{Mov}}\,(M) \ --- (**) \ .$$

Let $d \in \overline{\mathrm{Mov}}\,(M)^\circ$. Then d is represented by an effective $\mathbb{R}$-divisor on M, say D. By Kodaira's lemma, D is big. Choose a small positive real number $\epsilon > 0$ such that

$$(M, \epsilon D) = (M, K_M + \epsilon D)$$

is klt. As D is big, we can run the minimal model program for $(M, \epsilon D)$ to make D nef by [BCHM10, Theorem 1.2]. By Lemma 3.4, M has no divisorial contraction and all log-flips of $(M, \epsilon D)$ are τ_i $(i = 1, 2)$, as all log-flips of $(M, \epsilon D)$ are necessarily flops of M by $K_M = 0$. Thus, there is $g \in \mathrm{Bir}\,(M)$ such that $g^*D \in \overline{\mathrm{Amp}}\,(M)$. Therefore

$$\overline{\mathrm{Mov}}\,(M)^\circ \subset \mathrm{Bir}\,(M)^*\overline{\mathrm{Amp}}\,(M) = V^\circ \ --- (***) \ .$$

As both V and $\overline{\mathrm{Mov}}\,(M)$ are closed convex cone, (**) and (***) imply the result. □

We are now ready to complete the proof of Theorem 3.1.

By Lemma 3.2, we see that $d_1(f) = (2n^2-1) + 2n\sqrt{n^2-1} > 1$ and $f^*|N^1(M)$ is irreducible over $\mathbb{Q}$. Let $D \neq 0$ be a movable divisor on M. Then, as the class of D is rational and the both boundary rays of $\overline{\mathrm{Mov}}\,(M)$ is irrational by the first part of Lemma 3.5, the class of D belongs to $\overline{\mathrm{Mov}}\,(M)^\circ$. Thus by the second part of Lemma 3.5, there is $g \in \mathrm{Bir}\,(M)$ such that $g^*D \in \overline{\mathrm{Amp}}\,(M)$. As remarked at the beginning of this section, every nef divisor on M is semi-ample. In particular, g^*D is semi-ample. Now, we can apply Theorem 1.2 to conclude. □

References

[AC13] Amerik, E., Campana, F., : *Fibrations Méromorphes Sur Certaines Variétés à Fibré Canonique Trivial,* Pure and Applied Mathematics Quarterly, Special Issue: In honor of Fedor Bogomolov, **4** (2008) 509–545.

[AV16] Amerik, E., Verbitsky, M., : *Construction of automorphisms of hyperkähler manifolds*, arXiv:1604.03079.

[BCHM10] Birkar, C., Cascini, P., Hacon, C. D.; McKernan, J., : *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010) 405–468.

[Bi16] Bianco, F. L., : *On the primitivity of birational transformations of irreducible symplectic manifolds*, arXiv:1604.05261.

[Br67] Birkhoff,G., : *Linear transformations with invariant cones*. Amer. Math. Monthly **74** (1967) 274–276.

[Bo91] Borcea, C., : *Homogeneous vector bundles and families of Calabi-Yau threefolds. II.*, Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989), 83–91, Proc. Sympos. Pure Math. **52** Part 2, Amer. Math. Soc., Providence, RI, 1991.

[DS05] Dinh, T.-C., Sibony, N., : *Une borne supérieure de l'entropie topologique d'une application rationnelle,* Ann. of Math., **161** (2005) 1637–1644.

[Fu84] Fulton, W., : *Intersection Theory*, A Series of Modern Surveys in Mathematics, , Springer-Verlag, 1984.

[Hi64] Hironaka, H., : *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II.*, Ann. of Math. **79** (1964) 109–203; **79** (1964) 205–326.

[Ka88] Kawamata, Y., : *Crepant blowing-up of* 3*-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. of Math. **127** (1988) 93–163.

[Ka08] Kawamata, Y., : *Flops connect minimal models*, Publ. Res. Inst. Math. Sci. **44** (2008) 419–423.

[Ka09] Kawamata, Y., : *Finite generation of a canonical ring*, Current Developments in Mathematics 2007, International Press (2009) 43–76, arXiv:0804.3151.

[Og14] Oguiso, K., : *Automorphism groups of Calabi-Yau manifolds of Picard number 2*, J. Algebraic Geometry **23** (2014) 775–795.

[Og15] Oguiso, K., : *Some aspects of explicit birational geometry inspired by complex dynamics*, Proceedings of the International Congress of Mathematicians, Seoul 2014 (Invited Lectures) Vol.II (2015) 695–721.

[Og16-1] Oguiso, K., : *Simple abelian varieties and primitive automorphisms of null entropy of surfaces* : in K3 Surfaces and their Moduli, Progress in Math., Birkhäuser Progress in Mathematics **315** (2016) 279–296.

[Og16-2] Oguiso, K., : *Pisot units, Salem numbers and higher dimensional projective manifolds with primitive automorphisms of positive entropy*, arXiv:1608.03122.

[Ou16] Ouchi, G., : *Automorphisms of positive entropy on some hyperKähler manifolds via derived automorphisms of K3 surfaces*, arXiv:1608.05627

[Se06] Sernesi, E., : *Deformation of algebraic schemes*, Grundlehren der Mathematischen Wissenschaften, **334**, Springer-Verlag, 2006.

[Tr15] Truong, T.T., : *(Relative) dynamical degrees of rational maps over an algebraic closed field*, arXiv:1501.01523.

[Ue75] Ueno, K., *Classification theory of algebraic varieties and compact complex spaces,* Lecture Notes in Mathematics, **439**, Springer-Verlag, 1975.

[Zh09] Zhang, D.-Q., : *Dynamics of automorphisms on projective complex manifolds,* J. Differential Geom. **82** (2009) 691–722.

Mathematical Sciences, the University of Tokyo, Meguro Komaba 3-8-1, Tokyo, Japan and Korea Institute for Advanced Study, Hoegiro 87, Seoul, 133-722, Korea

E-mail address: oguiso@ms.u-tokyo.ac.jp